

Dissipation peak as an indicator of sample inhomogeneity in solid ^4He oscillator experiments

David A. Huse and Zuhair U. Khandker

Department of Physics, Princeton University, Princeton, NJ 08544

A simple phenomenological model is developed for the recent torsional oscillator experiments on solid ^4He . Within this model, for a homogeneous sample there is a specific quantitative relation between the change in the oscillator's frequency and its maximum damping at the apparent supersolid transition. Much of the published data do not satisfy this relation, indicating that the dissipation peaks in those samples are strongly inhomogeneously broadened.

PACS numbers: 67.80.-s, 67.80.Mg

Kim and Chan [1, 2, 3], and more recently Rittner and Reppy [4, 5], report apparent superfluidity in torsional oscillator experiments on *solid* ^4He . They freeze ^4He within a high- Q oscillator and measure the period and the damping of the oscillator as a function of the temperature T . As T is decreased, in a temperature range near 50-200 mK the oscillator's period is observed to decrease, indicating a reduction of the effective moment of inertia of the oscillator at low T . This is interpreted as the development of superfluidity within the solid, with the superfluid component remaining stationary and thus no longer contributing to the moment of inertia of the oscillator at low T . In the temperature range where the oscillator's period is most strongly temperature-dependent, the oscillator's damping has a maximum (equivalently, a minimum in the quality factor Q of the oscillator's resonance).

A simple phenomenological model of these experiments is of a two-component compound torsional oscillator:

$$(I - I_s) \frac{d^2\theta}{dt^2} = -G\theta - \Gamma(T) I_s \frac{d(\theta - \phi)}{dt}$$

$$I_s \frac{d^2\phi}{dt^2} = -\Gamma(T) I_s \frac{d(\phi - \theta)}{dt}.$$

The *relative* angular motion of the two components, $d(\phi - \theta)/dt$, is damped at a rate $\Gamma(T)$ that is strongly T -dependent. This relative motion experiences no restoring force. This motion represents the superfluid component (ϕ) moving with respect to the "normal" solid (θ), with the damping likely coming from the dissipative motion of thermally-excited vortices in the superfluid. At high T the damping is strong so $\Gamma(T) \gg \omega \approx \sqrt{G/I}$, where ω is the resonant frequency. The two components are then effectively locked together with total moment of inertia I . At low T the damping is weak or zero, $\Gamma(T) \ll \omega$, so the smaller, nominally superfluid component, with moment of inertia $I_s \ll I$ and angle ϕ , remains essentially at rest (the T -dependence of I_s is assumed to be much weaker than that of Γ , and thus is ignored). The resulting fractional change in the resonant frequency from high T to low T is $\frac{\Delta\omega}{\omega} \approx \frac{I_s}{2I} \ll 1$, due to the decrease of the moving moment of inertia from I to $I - I_s$. As

$\Gamma(T)$ passes through ω , this simple model has a dissipation peak, with the superfluid component then moving substantially both in the lab frame and relative to the solid. Let the magnitude of this damping peak be measured by $\Delta(Q^{-1})$, the increase in the oscillator's inverse quality factor Q^{-1} at this peak. Our simple model gives $\Delta(Q^{-1}) \approx \frac{I_s}{2I}$.

This simple result, $\Delta(Q^{-1}) \cong \frac{\Delta\omega}{\omega}$, is roughly consistent with torsional oscillator experiments on superfluid *liquid* ^4He films by Bishop and Reppy [6]. This result was also obtained from a more realistic model of superfluid liquid films within a Kosterlitz-Thouless theory by Ambegaokar *et al.* [7].

However, the ratio of the magnitude of the dissipation peak $\Delta(Q^{-1})$ to the fractional frequency shift $\frac{\Delta\omega}{\omega}$ varies widely among the published data on *solid* ^4He oscillator experiments [2, 3, 4, 5]. This is likely due to inhomogeneous broadening of the dissipation peak, from local "decoupling" transitions occurring at different temperatures in different parts of a macroscopically inhomogeneous polycrystalline solid, with the superfluidity arising on a network of grain boundaries or other defects in the solid [8, 9]. The total frequency shift $\Delta\omega$ from high to low T is set by the total amount of moment of inertia that decouples from the oscillator. For strong inhomogeneity, the T range over which this happens reflects the variation of the local transition temperatures. The damping at temperature T is only due to those parts of the solid that are passing through their transition at that T so that they have a local $\Gamma(T)$ that is near ω . Thus the magnitude of the dissipation peak $\Delta(Q^{-1})$ in an inhomogeneous solid is reduced from $\frac{\Delta\omega}{\omega}$ by the fraction of the sample that is passing through its local decoupling transition and thus contributing to the damping at the dissipation peak.

By this criterion, the data of Kim and Chan [2, 3] suggest they had much more inhomogeneous samples than those of the first paper of Rittner and Reppy [4]. Among those published, the result of Kim and Chan that appears the most homogeneous is that for pressure 30 bar shown in Fig. 2 of Ref. [3] (we restrict our attention to the lowest amplitude of oscillation). This run shows $\Delta(Q^{-1}) \cong 3 \times 10^{-6}$, about a factor of 6 or 7 smaller than its $\frac{\Delta\omega}{\omega} \cong 20 \times 10^{-6}$. In other examples

from Kim and Chan [2, 3] this factor ranges up to about 100. In their first paper, Rittner and Reppy [4], on the other hand, show data closer to the homogeneous expectation of $\Delta(Q^{-1}) \cong \frac{\Delta\omega}{\omega}$. E.g., their Fig. 2 shows $\Delta(Q^{-1}) \cong 12 \times 10^{-6}$ and $\frac{\Delta\omega}{\omega} \cong 18 \times 10^{-6}$. (The sample they label “First Run” even has $\Delta(Q^{-1})$ roughly 3 times *larger* than $\frac{\Delta\omega}{\omega}$, which is peculiar.) In Rittner and Reppy’s more recent paper [5], they intentionally made quite disordered samples, and these samples do show much weaker dissipation peaks indicating they are also quite inhomogeneous, as might be expected.

Another important difference between the published

results from these two labs is that for Rittner and Reppy [4] the transition disappeared in one crystal after annealing, while Kim and Chan [3] saw no large changes due to any annealing of their samples. This difference might also be related to differences in the sample inhomogeneity. One possible cause of sample inhomogeneities that would survive annealing is stresses due to temperature gradients that are present in the sample cell during the annealing and/or during the post-anneal cooling.

We thank Phil Anderson, Bill Brinkman, Moses Chan, Tony Clark, and John Reppy for discussions. Support was from the NSF through MRSEC grant DMR-0213706.

-
- [1] E. Kim and M. H. W. Chan, *Nature* **427**, 225 (2004).
 - [2] E. Kim and M. H. W. Chan, *Science* **305**, 1941 (2004).
 - [3] E. Kim and M. H. W. Chan, *Phys. Rev. Lett.* **97**, 115302 (2006).
 - [4] A. S. C. Rittner and J. D. Reppy, *Phys. Rev. Lett.* **97**, 165301 (2006).
 - [5] A. S. C. Rittner and J. D. Reppy, *cond-mat/0702665*.
 - [6] D. J. Bishop and J. D. Reppy, *Phys. Rev. Lett.* **40**, 1727 (1978); *Phys. Rev. B* **22**, 5171 (1980).
 - [7] V. Ambegaokar, B. I. Halperin, D. R. Nelson and E. D. Siggia, *Phys. Rev. Lett.* **40**, 783 (1978); *Phys. Rev. B* **21**, 1806 (1980).
 - [8] N. V. Prokof’ev and B. V. Svistunov, *Phys. Rev. Lett.* **94**, 155302 (2005).
 - [9] L. Pollet, M. Boninsegni, A. B. Kuklov, N. V. Prokof’ev, B. V. Svistunov and M. Troyer, *cond-mat/0702159*.